Does season of burn influence fuel life cycle and fire behavior in longleaf pine flatwoods of the southeastern US?



Jim Cronan, PhD student

University of Washington, School of Environmental and Forest Sciences

Overview

- Background
- Hypothesis
- Methods
- Results
- Conclusions
- Further analysis

Background: fuel treatments



Prescribed fire, St. Marks National Wildlife Refuge, Florida. J. Cronan, 2010.

Background: the longleaf pine ecosystem




Longleaf pine flatwoods. Apalachicola National Forest, Florida. J. Cronan, 2012.

Background: reference conditions



Pre-fire suppression flatwoods, southern Florida. John K Small, 1928.

Background: reference conditions



A. Annual Summer Burn

Waldrop et al. 1987


Hypotheses

 H₁: Postfire regeneration of herbaceous biomass is not equal between treatments.

 H₂: Postfire regeneration of woody biomass is not equal between treatments.


 H₃: Fire behavior (i.e. rate of spread, flame length, and reaction intensity) is not equal between treatments.

Methods: study area




Methods: sampling design


- Systematic plot layout fit to natural stand boundaries
- Fuel measures necessary to develop FCCS fuelbed.




Results: understory vegetation – departure from prefire loading




Results: Vegetation interaction plots




Results: FCCS fire behavior outputs




Results: FCCS output interaction plots




Results: mean peak fire temperature by site and burn date



Results: departure from prefire loading distribution by season and region



Results: NMDS plot, more mesic species at Eglin?



Conclusions

- Fail to reject null hypotheses. No effect of season on live fuel loading or predicted fire behavior.
- Mildly significant interaction between season and sampling episode.
- Inverse relationship between temperature and shrub regeneration.



Further Analysis

- Quantify relationship between burn temperature profiles and vegetation response.
- Impact of short-term (1990-2010) fire history and other ecological drivers.



Acknowledgements

- •Committee members
 Ernesto Alvarado, Clint Wright, John Skalski,
 Morgan Varner, Mark Kot
- Pacific Wildland Fire Sciences Lab
 Maria Petrova , Jon Dvorak, Jon McDuffey,
 Travis Freed, Cameron Balog, Aarin
 Sengsirirak, Paige Eagle, Bob Vihnanek, and
 Maureen Kennedy
- Funding provided by:



Supporting organizations

Eglin AFB: J. Kevin Hiers and Brett Williams.

Apalachicola NF: Eugene Watkins.

St. Marks NWR: Jennifer Hinckley and

Gregory Titus.





Results: rANOVA for vegetation

Herbaceous

	SS	df	SS _{error}	df	F statistic	Р
Season	0.0023	1	1.5748	7	0.0101	0.92283
Episode	4.4779	2	1.0871	14	28.8340	0***
Season:Episode	0.1403	2	1.7616	14	0.5577	0.58476
Woody						
Season	2.641	1	20.677	7	0.8942	0.37585
Episode	33.399	2	7.187	14	32.5293	0***
Season:Episode	0.424	2	1.796	14	1.6542	0.22652

Significance codes: ``***' < 0.001, `**' < 0.01, `*' < 0.05, ``' < 0.1.

- Bartlett's test for constant variance.
- Residuals normally distributed (Shapiro-Wilkes Normality test).

Results: rANOVA for FCCS outputs

Rate of Spread

	SS	df	SS _{error}	df	F	Р		
Season	0.177	1	0.597	7	2.0766	0.192764		
Episode	0.689	2	0.449	14	10.7326	0.001494**		
Season:Episode	0.358	2	0.433	14	5.7896	0.01712*		
Flame Length								
Season	0.005	1	3.93	7	0.0855	0.778479		
Episode	0.083	2	0.088	14	6.5464	0.009838**		
Season:Episode	0.039	2	0.073	14	3.7340	0.050159.		
Reaction Intensity								
Season	945	1	1045847	7	0.0063	0.9388		
Episode	18517	2	118813	14	1.0909	0.3628		
Season:Episode	22072	2	103062	14	1.4991	0.2571		

Results: shrub loading distribution by season and region

